

SCIENCE

Published by the **American Association for the Advancement of Science (AAAS)**, *Science* serves its readers as a forum for the presentation and discussion of important issues related to the advancement of science, including the presentation of minority or conflicting points of view, rather than by publishing only material on which a consensus has been reached. Accordingly, all articles published in *Science*—including editorials, news and comment, and book reviews—are signed and reflect the individual views of the authors and not official points of view adopted by the AAAS or the institutions with which the authors are affiliated.

The American Association for the Advancement of Science was founded in 1848 and incorporated in 1874. Its objectives are to further the work of scientists, to facilitate cooperation among them, to foster scientific freedom and responsibility, to improve the effectiveness of science in the promotion of human welfare, to advance education in science, and to increase public understanding and appreciation of the importance and promise of the methods of science in human progress.

Membership/Circulation

Director: Michael Spinella
Deputy Director: Marlene Zendell
Member Services: Rebecca Dickerson, *Manager*; Mary Curry, *Supervisor*; Pat Butler, Helen Williams, Laurie Baker, *Representatives*
Marketing: Dee Valencia, *Manager*; Jane Pennington, *Europe Manager*; Hilary Baar, *Associate*; Angela Mumeka, *Coordinator*
Research: Renuka Chander, *Manager*
Business and Finance: Jacquelyn Roberts, *Manager*
Administrative Assistant: Nina Araujo de Kobes
Science Member Services
 Marion, Ohio: 800-347-6969;
 Washington, DC: 202-326-6417
Other AAAS Programs: 202-326-6400

Advertising and Finance

Associate Publisher: Beth Rosner
Advertising Sales Manager: Susan A. Meredith
Recruitment Advertising Manager: Janis Crowley
Business Manager: Deborah Rivera-Vienhold
Finance: Randy Yi, *Senior Analyst*; Shawn Williams, *Analyst*
Marketing: John Meyers, *Manager*; Allison Pritchard, *Associate*
Traffic Manager: Carol Maddox
Recruitment: Terri Seiter, *Assistant Manager*; Pamela Sams, *Production Associate*; Debbie Cummings, Celeste Miller, Rachael Wilson, *Sales*; Charlotte Longhurst, *European Sales*
Reprints Manager: Corrine Harris
Permissions Manager: Lincoln Richman
Exhibits Coordinator: Arlene Ennis

PRODUCT ADVERTISING SALES: East Coast/E.

Canada: Richard Teeling, 201-904-9774, FAX 201-904-9701 • **Midwest/Southeast:** Elizabeth Mosko, 312-665-1150, FAX 312-665-2129 • **West Coast/W. Canada:** Neil Boylan, 415-673-9265, FAX 415-673-9267 • **UK, Scandinavia, France, Italy, Belgium, Netherlands:** Andrew Davies, (44) 1-457-838-519, FAX (44) 1-457-838-898 • **Germany/Switzerland/Austria:** Tracey Peers, (44) 1-270-760-108, FAX (44) 1-270-759-597 • **Japan:** Mashy Yoshikawa, (3) 3235-5961, FAX (3) 3235-5852
RECRUITMENT ADVERTISING SALES: US: 202-326-6555, FAX 202-682-0816 • **Europe:** Gordon Clark, (44) 1-81539-5211, FAX (44) 1223-302068 • **Australia/New Zealand:** Keith Sandell, (61) 02-922-2977, FAX (61) 02-922-1100

Send materials to *Science* Advertising, 1333 H Street, NW, Washington, DC 20005.

Information for Contributors appears on pages 112–114 of the 6 January 1995 issue. Editorial correspondence, including requests for permission to reprint and reprint orders, should be sent to 1333 H Street, NW, Washington, DC 20005.
Internet addresses: science_editors@aaas.org (for general editorial queries); science_letters@aaas.org (for letters to the editor); science_reviews@aaas.org (for returning manuscript reviews); membership@aaas.org (for member services); science_classifieds@aaas.org (for submitting classified advertisements); science_advertising@aaas.org (for product advertising)

LETTERS

Soil Erosion Estimates and Costs

In their article "Environmental and economic costs of soil erosion and conservation benefits," David Pimentel *et al.* (Articles, 24 Feb., p. 1117) assert that soil erosion is a major threat to the sustainability of agriculture all around the world and more specifically in the United States. In the source (1) they cite to support their introductory statement, Lal and Stewart (2) state that annual global erosion is about 36 billion tons, 10 billion attributable to natural causes and 26 billion to human activity. Lal and Stewart, in turn, cite a paper by Brown (3) as source for the 26-ton figure. In a review of Brown's work, I found (4) that his estimate of global erosion is based mainly on erosion estimates in the United States and on an extrapolation of the U.S. experience to the rest of the world. I did not and do not claim that Brown's estimate is wrong but the estimate rests on such thin underpinnings that it cannot be taken seriously.

Until the publication of work by Dregne and Chou (5) and Oldeman *et al.* (6) in the early 1990s, none of whom are cited by Pimentel *et al.*, there were no reliable estimates of how much erosion is occurring around the world, let alone its productivity consequences (2, 7–9).

The study by Dregne and Chou (5) deals with global degradation of rainfed cropland, irrigated land, and rangeland in dry areas, meaning arid, semi-arid, and dry subhumid climatic zones. Dregne and Chou classified these lands as slightly degraded (0 to 10% loss of productivity), moderately degraded (10 to 25% loss), severely degraded (25 to 50% loss), and very severely degraded (greater than 50% loss). I used their data (5) to calculate a weighted average degradation-induced loss of productivity of 11%. This is the cumulative loss over some period of time, which Dregne and Chou do not indicate. But for most of this land the period must be not less than several decades. The annual rate of productivity loss, therefore, would be less than 0.5%.

Oldeman *et al.* (6) found 1.965 billion hectares of degraded land around the world,

85% of it attributable to water and wind erosion. Thirty-eight percent of the degraded land was lightly degraded, 48% was moderately degraded, and 14% was strongly (or extremely) degraded. Oldeman *et al.* did not assign percentages of productivity loss to their degree-of-degradation categories. I assumed that the percentages correspond to those in the study by Dregne and Chou (5), that is, lightly degraded land has lost 0 to 10% of its productivity, and so on; I used the data from the study by Oldeman *et al.* to calculate a weighted average loss of 17%. The estimates of Oldeman *et al.* specifically refer to human-induced land degradation that occurred between the end of World War II and about 1990. The cumulative productivity loss of 17% over this 45 years implies an average annual loss of 0.4%.

Pimentel *et al.* cite a paper by Speth (10) as the source for the statement that "About 80% of the world's agricultural land suffers moderate to severe erosion, and 10% suffers slight to moderate erosion." However, Oldeman *et al.* (6) show that on a global scale about 1.03 billion hectares of agricultural land have suffered moderate-to-strong degradation because of wind and water erosion. This is less than 25% of the roughly 4.5 billion hectares of land in crops, pasture, and range

around the world, and well under a third of the 80% figure given by Pimentel *et al.*

Pimentel *et al.* cite a paper by Barrow (11) for their assertion that soil erosion rates are highest in Asia, South America, and Africa, averaging 30 to 40 tons per hectare per year. Barrow states (11, p. 209) that the estimates he discusses are crude and that it "probably would be wise to wait until the publication of the GLASOD [Global Assessment of Soil Degradation] maps, sometime after 1990, before trying to get an accurate overview of soil erosion." The GLASOD maps are those prepared by Oldeman *et al.*, published in 1990 (6).

Pimentel *et al.* state that over the last 200 years 100 million hectares (about 30%) of U.S. farmland has been abandoned because of erosion, salinization, and waterlogging and cite the U.S. Department of Agriculture



Soil erosion. Corn field in Missouri in 1994 shows effects of cultivation.

J.P. JACKSON/PHOTO RESEARCHERS

(USDA) (12), Bennett (13), and Pimentel *et al.* (14). In a close reading of the USDA report (12), I found nothing about degradation of U.S. land over the last 200 years, and the Bennett citation is a 1939 publication.

Pimentel *et al.* cite the USDA (12) as the source for the statement that the combined effect of water and wind erosion moves an average of 17 tons per hectare per year from U.S. croplands, which figure they then use in cost estimates. Successive USDA surveys (15) provide more accurate estimates of cropland erosion for 1982, 1987, and 1992. Pimentel *et al.* do not reference the updated surveys which show that the 1992 rate was 13 tons per hectare per year, almost 25% less than originally reported (12).

Pimentel *et al.* state that they have "developed empirical models that incorporate the numerous factors affecting both erosion rates and soil productivity." However, the models are not presented, so the reader is



Water erosion. Farm field washing away in Harper County, Kansas, in 1984.

LARRY MILLER/PHOTO RESEARCHERS

unable to evaluate them. We are told that the models are based on numerical assumptions about rainfall, soil depth, type and slope of soil, percent of soil organic matter, and an annual erosion rate of 17 tons per hectare per year. No sources are given for any of these numbers, although the last evidently is from (12) and hence is for 1982. The numbers, and presumably the unspecified models, are used to estimate the annual on-farm per hectare economic costs of losses of soil and water resulting from erosion of 17 tons per hectare

per year on conventionally tilled land in corn, over a 20-year period (Pimentel *et al.*'s table 2). This estimate is then multiplied by 160 million hectares, said to be the total amount of cropland in the country, to get an estimate of the annual nationwide on-farm economic costs of cropland erosion. This estimate is \$27 billion per year, although the per hectare estimate of on-farm costs of \$146 (Pimentel *et al.*'s table 2), when multiplied


by 160 million hectares, gives a total cost of \$23 billion (not \$27 billion).

These procedures, the numbers used, and the results obtained prompt several questions and comments.

1) Why should the assumed conditions with respect to precipitation, soil type, slope, depth, and percent organic matter be representative of cropland in the country as a whole? These conditions are highly variable across regions.

2) How was the \$100 per hectare cost of nutrient replacement estimated (Pimentel *et al.*'s table 2)? A source is cited for the losses of nutrients in terms of kilograms, but no information is given about how these losses were valued. The issue is of major importance because nutrient losses account for two-thirds of the total on-farm economic costs. Multiplying average 1992 prices for anhydrous ammonia, the most common form of nitrogen fertilizer, superphosphate (44 to 46% phosphate), and potassium chloride (60% potassium) (16, p. 27) by the quantities of lost nutrients shown by Pimentel *et al.* (Pimentel *et al.*'s table 3) gives an estimate of plant-available nutrient losses of about \$23 per hectare. Even if the cost is measured by total nutrient losses, that is, by counting nutrients not available to support plant growth in any given year, the total per hect-

This is my system for affinity separations of polyclonal and monoclonal antibodies, enzymes and fusion proteins.



are cost of nutrient losses comes to only about \$75, still well below \$100.

3) What are we to make of the cost of the erosion-induced losses of water (Pimentel *et al.*'s table 2)? At \$30 per hectare per year, this cost is 20% of total on-farm costs. A note in that table says that it is the cost of supplying groundwater for irrigation to replace erosion-induced losses of water from precipitation: "if rainfall were abundant, then this replacement cost would not be necessary." In the main crop-producing areas of the country east of the Great Plains, rainfall is generally adequate to maintain current yields, as indicated by the scant use of irrigation. In those areas the cost of replacing erosion-induced losses of water, as estimated by Pimentel *et al.*, should be zero. Pimentel *et al.* acknowledge this problem, but include the estimated cost of water losses in their calculation of nationwide costs.

Other studies show much lower on-farm costs of soil erosion in the United States than Pimentel *et al.* do. One such study is based on the Erosion Productivity Impact Calculator (EPIC) model developed by USDA soil scientists (12). EPIC simulates the productivity effects of soil erosion on soil characteristics and processes, including losses of soil nutrients, water-holding capacity, and acidity (pH). In (12), the estimate

with EPIC showed the average annual gross on-farm costs of 100 years of cropland erosion in the United States at 1982 rates to be \$252 million. In another study (17), I used results from the Productivity Index model, developed by soil scientists at the University of Minnesota (18), to estimate the annual gross cost of erosion-induced on-farm losses of productivity in the United States to be \$500 million to \$600 million.

The present rate of cropland erosion in the United States is probably close to 13 tons per hectare per year, not 17. Pimentel *et al.* have greatly overestimated the on-farm per hectare costs of replacing nutrients and water. Their estimate of the nationwide on-farm costs of cropland erosion appears to be greatly overstated, even if their procedures and assumptions are accepted.

Pierre Crosson

Resources for the Future,
1616 P Street, NW,
Washington, DC 20036, USA

References and Notes

1. N. Myers, *Gaia: An Atlas of Planet Management* (Anchor and Doubleday, Garden City, NY, 1993).
2. R. Lal, in *Soil Degradation*, R. Lal and B. A. Stewart, Eds. (Springer-Verlag, New York, 1990), p. 132.
3. L. Brown, *State of the World 1984* (Worldwatch Institute, W. W. Norton, New York, 1984).
4. P. Crosson, in *Population and Food in the Early 21st Century: Meeting Future Food Demands of Increasing Population*, N. Islam, Ed. (International Food Policy Research Institute, Washington, DC, 1995), pp. 143–159.
5. H. E. Dregne and Nan-Ting Chou, in *Degradation and Restoration of Arid Lands* (Texas Tech Univ., Lubbock, TX, 1992), pp. 249–282.
6. L. Oldeman, R. Hakkeling, W. Sombroek, *World Map of the Status of Human-Induced Soil Degradation: An Explanatory Note* (International Soil Reference and Information Center, Wageningen, The Netherlands, and United Nations Environment Programme, Nairobi, Kenya, 1990).
7. R. Nelson, *Dryland Management: The Land Degradation Problem* (Environment Department Working Paper No. 8, World Bank, Washington, DC, 1988).
8. H. E. Dregne, in *Challenges in Dryland Agriculture: A Global Perspective*, P. Unger, T. Sneed, W. Jordan, R. Jensen, Eds. (Texas Agricultural Experiment Station, Amarillo/Bushland, TX, 1988), pp. 610–612.
9. S. El-Swaify, E. Dangler, C. Armstrong, *Soil Erosion by Water in the Tropics* (Univ. of Hawaii, Honolulu, 1982).
10. J. G. Speth, *Towards an Effective and Operational International Convention on Desertification* (International Negotiating Committee, International Convention on Desertification, United Nations, New York, 1994).
11. C. J. Barrow, *Land Degradation* (Cambridge Univ. Press, Cambridge, UK, 1991).
12. *The Second RCA Appraisal: Soil, Water, and Related Resources on Nonfederal Land in the United States* (U.S. Department of Agriculture, Washington, DC, 1989).
13. H. H. Bennett, *Soil Conservation* (McGraw-Hill, New York, 1939).
14. D. Pimentel *et al.*, *Science* **194**, 149 (1976).
15. *Summary Report 1992 National Resources Inventory* (Soil Conservation Service, U.S. Department of Agriculture, Washington, DC, 1994).
16. H. H. Taylor, *Fertilizer Use and Price Statistics, 1960–93* (Statistical Bulletin No. 893, Economic Re-

Yeah right, where's the rest of it?

Can you put a syringe and a HiTrap® together? Well, that's all it takes to master your system for doing affinity separations in molecular biology.

While HiTrap offers a nearly effortless solution to setting up your affinity separation system, it's also proficient at saving you time and money. That's because the affinity media pre-packed into HiTrap run at high flow rates and high sample loadings. You'll get high recovery separations and quick results—in just a few minutes you can purify polyclonal and mono-

clonal antibodies, enzymes and fusion proteins. HiTrap—the widest range of affordable affinity separations available for molecular biologists—can be delivered in 1 or 5 ml sizes.

Just call us at 1 (800) 526 3593 in the United States or +46 18 16 5011 from the rest of the world to find out more about HiTrap and its effortless affinity separation procedure.

We probably shouldn't say procedure, though. After all, anything so fast and easy can't really be called a procedure.



**Pharmacia
Biotech**

Uppsala, Sweden. (And the rest of the world.)

search Service, U.S. Department of Agriculture, Washington, DC, 1994).

17. P. Crosson, in *Agriculture and the Environment*, T. Phipps, P. Crosson, K. Price, Eds. (Resources for the Future, Washington, DC, 1986), pp. 35-73.
18. F. Pierce, R. Dowdy, W. Larson, W. Graham, *J. Soil Water Conserv.* **39**, 131 (1984).

Response: Crosson indicates that the estimate of 75 billion metric tons per year of world soil lost to erosion, worldwide, is too high (1). The estimated soil loss in the United States is nearly 4 billion tons per year on nonfederal land (2) plus an estimated 0.5 billion tons on federal lands. The United States has about 11% of the world's arable land and approximately the same percentage of pasture land (3). Therefore, assuming that the rest of the world suffers similar rates of erosion, the total global soil loss would be approximately 40 billion tons per year. However, as pointed out in our report, soil erosion rates in Asia, Africa, and South America are about double those of the United States (4, 5). Taking these higher rates of erosion into consideration, the estimated 75 billion tons per year of eroded soil seems reasonable.

The estimate that 80% of the world's agricultural land suffers from moderate to severe soil degradation (6) is consistent with several other investigations (4, 5).

Bennett (7) in 1939 reported that about

80 million hectares of cropland either had been ruined, severely damaged, or had lost one-half of all topsoil. Since 1939, U.S. agricultural land has continued to erode and be lost to production (8). Our estimate, that 100 million hectares (about 30%) of U.S. farmland has been abandoned, is conservative. Lal (9) reports that an estimated 2.0 billion hectares of once productive agricultural land has been degraded or destroyed during the history of agriculture worldwide. Agricultural land continues to be degraded and abandoned because of erosion and is resulting in the rapid and continued spread of agriculture into world forest-lands (10).

We did not see the latest USDA (11) survey that was published in 1994 on soil erosion because our paper was submitted during the summer of 1994. We are delighted to know that during the past 10 years soil erosion on cropland has declined by 25%. However, the current erosion rate of 13 tons per hectare per year is still 13 times above the soil sustainability rate. Also, the rates of soil erosion on pastures and rangelands in the survey did not decline and remain a serious threat to these agricultural lands (11).

Crosson states that rainfall east of the Great Plains, including the "corn belt," is adequate for corn production. However, adequate rainfall is not the same as optimum.

Corn production even in the corn belt usually suffers from water shortages during the summer growing season (12). Thus, the increased water loss associated with soil erosion has a negative impact on corn yields.

We stated in our article explicitly how the \$27-billion-per-year estimated nationwide on-farm economic cost of cropland erosion was calculated. This was based on a \$20 billion replacement value for soil nutrients (8) and \$7 billion for loss of water and reduced soil depth. We stated in detail the assumptions and documented the sources for the field experimental data used in our tables 2 and 3. We agree with Crosson that soil type, precipitation, slope, soil depth, organic matter, and soil biota vary from field to field and region to region and all have an effect on erosion and crop productivity. This is the reason that we carefully stated the conditions and assumptions for the assessments included in our tables 2 and 3.


Crosson indicates that the \$27 billion on-farm economic costs that we estimated are too high. In his earlier paper (13), he estimated that the total annual cost of lost nutrients was \$500 million for U.S. agriculture. This is in stark contrast to the \$18 billion for 1980 (14) and \$20 billion for 1991 (8) estimates of soil nutrient losses reported by several soil scientists at Iowa

Porsche Power on a PC, or a Mac, or a Sun Workstation.

If you use electrophoresis in your research, chances are you're not taking full advantage of the technique by using image analysis—probably because the software you tried was either difficult to use or limited in what applications it could support.

Perform a wide variety of analytical tasks quickly and easily with Bio Image IQ™ Software. This menu-driven package is self-installing and self-tutoring. IQ software supports whole band 1-D, 2-D, blot analysis, and colony / plaque counting. Prepare reports, annotate images for publication, and compare data in your favorite spreadsheet and database programs—all without leaving your keyboard.

Take IQ software for a test drive. To receive your demo disk, call us or E mail sales@bioimage.com.



 **Bio Image**

No matter what computer you use, Bio Image IQ™ Software lets you take full advantage of electrophoresis by making image analysis powerful yet easy-to-use.

Bio Image USA
TEL: (313) 930-9900
FAX: (313) 930-0990

Bio Image France
TEL: (1) 48.63.24.25
FAX: (1) 48.63.72.99

Bio Image U.K.
TEL: (44) 1270-886192
FAX: (44) 1270-886100

Nihon Bio Image, Ltd.
TEL: (81) 3-3280-0990 / 0932
FAX: (81) 3-3280-0991