

Set #4 - for Thurs May 3

Read Ohanian Ch. 5Read Feynman Vol. III Ch. 1, Ch.2, Ch. 3**From Ohanian:****Ch. 5** Problems 17, 18, 27, 28, 34.**Ch. 5** Problems 31 (find the standard deviation for Δx)

1. Show that if $\Delta x \Delta p = \frac{1}{2} \hbar$, the minimum energy of a simple harmonic oscillator is $\frac{1}{2} \hbar \omega$. What is the minimum energy in joules for a mass of 10^{-2} kg oscillating on a spring of force constant 1.0 N/m?

2. A particle is on a table in a uniform gravitational field. The energy is $E = mgz + p^2/2m$, where $z = 0$ at the table. Classically, the minimum energy is $E = 0$. In quantum mechanics, because of the uncertainty principle, the particle cannot be at rest at a well-defined position. Assume then that the particle moves in a small range Δz above $z = 0$ and take the average height of the particle $\bar{z} = \frac{1}{2} \Delta z$ and apply the uncertainty principle in the form $\Delta z \Delta p \geq \hbar/2$.

a) Show that the average energy of the particle satisfies: $\bar{E} \geq \frac{1}{2} mg \Delta z + \frac{\hbar^2}{8m(\Delta z)^2}$.

b) Minimize \bar{E} as a function of Δz and find \bar{E}_{\min} and \bar{z}_{\min} (the value of \bar{z} that minimizes \bar{E} .)

3. A student drops a marble of mass m from a tower of height H . The student uses a very precise apparatus to aim the marble as well as possible so as to hit a particular crack in the sidewalk.

a) Show that the marble will, on average, miss the crack by a distance of order

$$d = \sqrt{\frac{\hbar}{m}} \left(\frac{H}{g} \right)^{1/4}$$

where h is the Planck's constant and g is the acceleration due to gravity. (This formula ignores factors of 2 and π .)

b) Assuming reasonable values of H and m , estimate the values of d . How important are these quantum-mechanical effects?

4. Consider a continuous distribution of wave numbers $A(k)$ and consider the wavepacket at $t = 0$:

$$\psi(x) = \int_{-\infty}^{\infty} A(k)e^{ikx} dk$$
$$A(k) = \begin{cases} A_0 & ; \quad |k - k_0| < \Delta k/2, \\ 0 & ; \quad |k - k_0| > \Delta k/2 \end{cases}$$

a) Determine the wavefunction at $t = 0$: $\psi(x, 0)$.

b) Determine the probability density function $|\psi(x, 0)|^2$.

c) Sketch a graph of $|\psi(x, 0)|^2$ and discuss your result.

5. From the uncertainty principle, estimate the minimum kinetic energy of an electron confined to a nucleus of size 5 fm (1 fm = 10^{-15} m). Note: This is a relativistic electron. Why?